

CLICK HERE TO JUMP TO USE DIRECTIONS



TRIMEC® 899 BROADLEAF HERBICIDE

Controls: Dandelion, Knotweed, Henbit, Chickweed, Plantain, Spurge, and many other broadleaf weeds including those listed on this label.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	30.56%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	8.17%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)	2.77%
INERT INGREDIENTS:	58.50%
TOTAL 100.00%	

This Product Contains:

2.38 lbs. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 25.38%
0.63 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 6.75%
0.21 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.30%
Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.



**KEEP OUT OF REACH OF CHILDREN
DANGER**

See below for Precautionary Statements and First Aid.

KEEP FROM FREEZING



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Engineering Control Statements:

Containers over 1 gallon and less than 5 gallons: Persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron.

Containers of 5 gallons or more: Do not open pour product from this container. A mechanical system (probe and pump or siphon) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID

If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

(cont. on next column)

FIRST AID (continued)

If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.	
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	

ENVIRONMENTAL HAZARDS: This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP-p has been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D and MCPP-p pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

**DIRECTIONS FOR USE
FOR USE ON RESIDENTIAL AND OTHER TURF SITES
EXCLUDING SOD FARMS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: For Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. **For Metal Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

USE PRECAUTIONS

Do not apply this product through any type of irrigation system. Avoid drift of spray mist to vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not spray on carpetgrass, dichondra nor on lawns or turf where desirable clovers are present. Do not apply this product to 'Floritam' St. Augustinegrass. Use only lawn type sprayers. Coarse sprays are less likely to drift. Use coarse spray droplets. Avoid fine mists. Do not spray roots of ornamentals and trees. Do not exceed specified dosages for any area. Be particularly careful within the dripline of trees and other ornamental species. Do not apply this product to warm season grasses during spring green-up which is the transition period between dormancy and active growth. Do not apply to newly seeded grasses until well established. Do not broadcast apply when air temperatures exceed 85°F; some injury may be expected with spot treatments when air temperatures exceed 85°F. Seed can be sown 3 to 4 weeks after application at recommended rate. After using this product, clean sprayer with soap or detergent and water, and rinse thoroughly before applying other pesticides. Failure to observe the above precautions may result in injury.

ORNAMENTAL LAWNS & TURF

Apply 2.3 to 4.0 pints of product per acre in sufficient water for adequate coverage. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing. Avoid applying during long excessively dry or hot periods unless adequate irrigation is available. Do not irrigate within 24 hours of application. Do not apply to newly seeded turf until after the second or third mowing. Grass seed can be sown 3 to 4 weeks after application. Do not use on dichondra and carpetgrass.

The maximum application rate to turf is 1.3 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

APPLICATION TYPE

Kentucky Bluegrass, Fescue and Perennial Ryegrass:

Professional Lawn Maintenance or Mature Weeds: Use 2.3 to 4.0 pints of product per acre. Or, use 0.84 to 1.5 fl. oz. of product per 1,000 sq. ft. with spray volumes of 0.5 to 5 gallons of spray solution per 1,000 sq. ft. The best time to control clover is in the fall. When making a spring application to control clover, the 4 pint per acre rate is recommended.

Controlled Droplet Applicator: Reduced rates must be used when grass is stressed from heat or drought. Do not use on St. Augustinegrass or centipedegrass. Add 2 pints TRIMEC® 899 to the Herbi container, then fill with 3 pints of water. Spray entire contents over 33,000 sq. ft. (approx. ¾ acre). Do not overlap between spray patterns.

Bermudagrass and Zoysiagrass:

Apply 3.0 to 4.0 pints of product per acre with spray volumes ranging from 20 to 220 gallons per acre. The equivalent application rates are 1.1 to 1.5 fl. oz. of product per 1,000 sq. ft. with spray volumes of 0.5-5.0 gallons of spray solution per 1,000 sq. ft. Some hybrid bermudagrass may be sensitive to this product. Contact your local extension service weed control specialist.

St. Augustinegrass:

Apply 2.3 pints of product in up to 150 gallons of water per acre. (1.2 fl. oz. in 5 gallons water per 1,500 sq. ft.). Do not spray St. Augustinegrass if stressed from heat, drought, etc. Slight turf yellowing should disappear after about one week.

Bentgrass:

On closely mowed bentgrass (putting and bowling greens) apply at a maximum rate of one fluid ounce of product in 5 gallons water per 1,500 sq. ft. — preferably in May or mid-August through September. Slight turf yellowing will disappear after about one week.

NOTE: Care should be taken to avoid overdosing bentgrass and St. Augustinegrass or injury may result. High spray volumes (i.e. one fluid ounce in 5 gallons water per 1,500 square feet) will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle rather than a multiple nozzle boom. Sweeping movements with a multiple nozzle boom will result in heavy local over-application and subsequent turf discoloration or injury.

TRIMEC® 899 BROADLEAF HERBICIDE CONTROLS

Annual fleabane	Compassplant	Groundsel	Pennsylvania smartweed	Velvetleaf (*buttonweed)
Aster, white heath	Curly dock	Hairy bittercress	Pennywort (*dollarweed)	Venice mallow
& white prairie	Dandelion	Hawkweed	Pepperweed	Veronica (*corn speedwell)
Bedstraw	Dayflower	Healall	Pigweed	Virginia buttonweed
Beggarticks	Deadnettle	Henbit	Pineappleweed	Virginia-creeper
Beggarweed, creeping	Dock	Horsenettle	Plantain	Western salsify
Bindweed	Dogfennel	Horseweed	Poison ivy	White clover (*Dutch clover,
Birdsfoot trefoil	Dovefoot geranium	Innocence (Blue-eyed Mary)	Poison oak	honeysuckle clover,
Black medic	English daisy	Jimsonweed	Prostrate knotweed	white trefoil & purplewort)
Broadleaf plantain	False dandelion	Kochia	(*knotweed)	Wild carrot
Buckhorn plantain	(*spotted catsear	Lambsquarters	Puncturevine	Wild garlic
Bull thistle	& common catsear)	Lawn burweed	Purple cudweed	Wild geranium
Burclover	Field bindweed	Lespedeza, common	Purslane	Wild lettuce
Burdock, common	(*morningglory	Mallow, common	Ragweed	Wild mustard
Buttercup, creeping	& creeping jenny)	Matchweed	Redweed	Wild onion
Carolina geranium	Field madder	Mouseear chickweed	Red sorrel (*sheep sorrel)	Wild strawberry
Carpetweed	Field oxeye-daisy	Mustard	Roundleaf greenbriar	Wild violet
Chickweed, common	(*creeping oxeye)	Nettle	Shepherdspurse	Yarrow
Chicory	Field pennycress	Old world diamond flower	Spotted spurge	Yellow rocket
Cinquefoil	Filaree, whitestem & redstem	Oxalis (*yellow woodsorrel &	Spurge	
Clover	Florida pusley	creeping woodsorrel)	Sunflower	
Cocklebur	Ground ivy	Parsley-piert	Thistle	*Synonyms

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

899 AP021404
EPA REG. NO. 2217-694

Manufactured By

An Employee-Owned Company
1217 West 12th Street
Kansas City, Missouri 64101
consumer.pbigordon.com